

CEFTRIAXONE

Other names: **Rocephin**®, **AFT brand**®



Important note: The following information is intended to add to, not replace, the knowledge and judgment of your doctor, pharmacist or other health care professional.

Why are you taking this medication

Ceftriaxone is a cephalosporin antibiotic that kills bacteria. It is effective against many (but not all) infections. This medicine may be used for serious infections, where other antibiotics would not work as well.

When to take this medication

Ceftriaxone does not work if it is taken orally (as a tablet or capsule). It only works if it is injected into muscle (IM) or a vein (IV).

Ceftriaxone may already be prepared by the Hospital Pharmacy or will be prepared by a District Nurse. The injection is contained in a bag and is designed to drip slowly into your catheter (this is called an infusion) or as a syringe. This should be administered exactly as you have been instructed at the hospital or clinic. This infusion should be given over at least 30 minutes. The syringe should be given over at least five minutes.

Your dose (how much and how often) depends on how bad your infection is and on your kidney and liver function.

Side effects

Some people are allergic to ceftriaxone: this would have showed up in the hospital or clinic.

Milder side effects include:

- nausea
- diarrhoea
- headache

Contact your backup service or see your doctor **immediately** if the following occur:

- skin rash, itching or fever
- unusual bleeding or bruising

Rarely, ceftriaxone can cause severe or long lasting diarrhoea which may start up to a month after you finish taking ceftriaxone. If you get troublesome diarrhoea, do not take any anti-diarrhoea medicine. Contact your backup service or your doctor straight away and report that you have had ceftriaxone recently: there is a specific treatment for this diarrhoea.

Other medication

Ceftriaxone may increase the effect of **warfarin** on your blood clotting making you more prone to bleeding.

Storage

Remember to keep your medication out of reach of children

Read this leaflet and discuss any questions you have about your medicines with your doctor or pharmacist.