Children's medication information leaflet

ANTIEPILEPTIC DOSING CHART



For children when changing therapy

Your child's antiepileptic medicine has been altered. This may mean that one or more medicines may slowly be withdrawn (stopped) while at the same time one or more medicines may slowly be introduced. This is to ensure that seizures are controlled while minimising side effects of the new medicine(s).

WEEK ONE

Medication	Medication Strength	START Date	Morning Dose	Midday Dose	Teatime Dose	Bedtime Dose	STOP Date

WEEK TWO

Medication	Medication Strength	START Date	Morning Dose	Midday Dose	Teatime Dose	Bedtime Dose	STOP Date

WEEK THREE

Medication	Medication Strength	START Date	Morning Dose	Midday Dose	Teatime Dose	Bedtime Dose	STOP Date

WEEK FOUR

Medication	Medication Strength	START Date	Morning Dose	Midday Dose	Teatime Dose	Bedtime Dose	STOP Date

WEEK FIVE

Medication	Medication Strength	START Date	Morning Dose	Midday Dose	Teatime Dose	Bedtime Dose	STOP Date

WEEK SIX

Medication	Medication Strength	START Date	Morning Dose	Midday Dose	Teatime Dose	Bedtime Dose	STOP Date

WEEK SEVEN

Medication	Medication Strength	START Date	Morning Dose	Midday Dose	Teatime Dose	Bedtime Dose	STOP Date

WEEK EIGHT

Medication	Medication Strength	START Date	Morning Dose	Midday Dose	Teatime Dose	Bedtime Dose	STOP Date

Read this leaflet and discuss any questions you have about your medicines with your doctor or pharmacist.